

Jordan-Hölder theorems for derived categories of derived discrete algebras

Yongyun Qin

Chinese Academy of Sciences, Beijing 100190, P.R. China.

E-mail: qinyongyun2006@126.com

Abstract

For any positive integer n , n -derived-simple derived discrete algebras are classified up to derived equivalence. Furthermore, the Jordan-Hölder theorems for all kinds of derived categories of derived discrete algebras are obtained.

Mathematics Subject Classification (2010): 16G10; 18E30

Keywords : derived discrete algebra; n -derived-simple algebra; n -composition-series; n -composition-factor; 2-truncated cycle algebra.

1 Introduction

Throughout this paper, k is an algebraically closed field and all algebras are finite dimensional associative k -algebras with identity. Recollements of triangulated categories were introduced by Beilinson, Bernstein and Deligne [6], and play an important role in algebraic geometry and representation theory. Recollements, more general n -recollements [17], of derived categories of algebras provide a reduction technique for some homological properties such as the finiteness of global dimension [34, 26, 3], the finiteness of finitistic dimension [22, 15], the finiteness of Hochschild dimension [16] and homological smoothness [17], and some homological invariants such as K -theory [32, 36, 29, 31, 14, 3], Hochschild homology and cyclic homology [25] and Hochschild cohomology [16]. Moreover, some homological conjectures such as the finitistic dimension conjecture [22, 15], the Hochschild homology dimension conjecture [17], the Cartan determinant conjecture [17], the Gorenstein

symmetry conjecture [17] and the (dual) vanishing conjecture [35], can be reduced to n -derived-simple algebras by n -recollements for appropriate $n \geq 1$. A recollement of derived categories of algebras is a short exact sequence of derived categories of algebras, and thus leads to the concepts of composition series and composition factors and Jordan-Hölder theorem analogous to those in group theory and module theory. Indeed, the composition factors in this context are the derived categories of derived simple algebras which were studied in [34, 21, 27, 28, 3]. Meanwhile, the Jordan-Hölder theorem of derived category was established for hereditary algebras [1] and later for piecewise hereditary algebras [2]. Nonetheless, this theorem was disproved for certain infinite dimensional algebras [13, 14], and later for a finite dimensional algebra [3]. Hence, the Jordan-Hölder theorem of the derived category of an algebra is a little bit subtle.

Derived discrete algebras were introduced by Vossieck [33], which are crucial in the Brauer-Thrall type theorems for derived category [18] and were explored in [8, 7, 11, 12, 19, 9]. In particular, Vossieck proved that a basic connected derived discrete algebra is isomorphic to either a piecewise hereditary algebra of Dynkin type or some one-cycle gentle algebra which does not satisfying the clock-condition [33]. The latter was further classified up to derived equivalence by Bobiński, Geiß and Skowroński [8]. Synthesizing the relevant results of Vossieck [33], Happel [20] and Bekkert-Merklen [5], all indecomposable objects in the bounded derived category of a derived discrete algebra are fairly clear.

In this paper, we will classify all n -derived-simple derived discrete algebras up to derived equivalence, and prove the Jordan-Hölder theorems for all kinds of derived categories of derived discrete algebras. More precisely, we will show that, for any positive integer n , a derived discrete algebra is n -derived-simple if and only if it is derived equivalent to the underlying field k or a 2-truncated cycle algebra. In order to obtain simultaneously the Jordan-Hölder theorems for unbounded, upper bounded, lower bounded, bounded derived categories of derived discrete algebras, we will introduce n -composition-series and n -composition-factors of the unbounded derived category of an algebra. Furthermore, we will prove that the unbounded derived category of any derived discrete algebra admits an n -composition-series whose n -composition-factors are independent on the choice of n -composition-series up to reordering and derived equivalence. In particular, the Jordan-Hölder theorem holds for the unbounded (resp. upper bounded, lower bounded, bounded) derived category of any derived discrete algebra.

This paper is organized as follows: In section 2, we will study the n -recollements of derived categories of derived discrete algebras. In Section 3, we will classify all n -derived-simple derived discrete algebras up to derived equivalence. In section 4, we will define the n -composition-series and n -composition-factors of the unbounded derived category of an algebra, and prove the Jordan-Hölder theorems for derived categories of derived discrete algebras.

2 n -recollements on derived discrete algebras


In this section, we will study the n -recollements of derived categories of derived discrete algebras.

Let A be an algebra. Denote by $\text{Mod}A$ the category of right A -modules, and by $\text{mod}A$ (resp. $\text{proj}A$) its full subcategory consisting of all finitely generated modules (resp. finitely generated projective modules). Denote by $\mathcal{D}(\text{Mod}A)$ or just $\mathcal{D}A$ the unbounded derived category of A . Denote by $\mathcal{D}^b(\text{mod}A)$ or just $\mathcal{D}^b(A)$ the bounded derived category of A . Denote by $K^b(\text{proj}A)$ the homotopy category of cochain complexes of objects in $\text{proj}A$. An object X in $\mathcal{D}A$ is said to be *compact* if $\text{Hom}_{\mathcal{D}A}(X, -)$ commutes with direct sums. It is well-known that up to isomorphism the objects in $K^b(\text{proj}A)$ are precisely all the compact objects in $\mathcal{D}A$ (Ref. [10]).

An algebra A is said to be *derived discrete* provided for every vector $\mathbf{d} = (d_p)_{p \in \mathbb{Z}} \in \mathbb{N}^{(\mathbb{Z})}$ there are only finitely many isomorphism classes of indecomposable objects X in $\mathcal{D}^b(A)$ of cohomology dimension vector $(\dim_k H^p(X))_{p \in \mathbb{Z}} = \mathbf{d}$ (cf. [33]). Some characterizations of derived discrete algebras in terms of (global) cohomological range and (global) cohomological length are provided in [18].

Vossieck proved in [33] that a basic connected derived discrete algebra is isomorphic to either a piecewise hereditary algebra of Dynkin type or a one-cycle gentle algebra not satisfying the clock-condition. The latter was shown by Bobiński-Geiß-Skowroński in [8] to be derived equivalent to a gentle algebra $\Lambda(r, s, t)$ with $1 \leq r \leq s$ and $t \geq 0$, which is the bound quiver algebra

[4] given by the quiver $Q(s, t) :=$



with the relations $\alpha_{s-1}\alpha_0, \alpha_{s-2}\alpha_{s-1}, \dots, \alpha_{s-r}\alpha_{s-r+1}$. Thus, up to derived equivalence, a derived discrete algebra can be decomposed as $\bigoplus_{p=1}^m A_p$ where $m \geq 1$ and $A_p = \Lambda(s_p, s_p, t_p)$ for $s_p \geq 1, t_p \geq 0, 1 \leq p \leq u$ and $0 \leq u \leq m$; $A_p = \Lambda(r_p, s_p, t_p)$ for $s_p > r_p \geq 1, t_p \geq 0, u+1 \leq p \leq v$ and $u \leq v \leq m$; A_p is a basic connected piecewise hereditary algebra of Dynkin type for $v+1 \leq p \leq m$. Obviously, the algebra $\Lambda(r, s, t)$ is of finite global dimension if and only if $s > r$. The algebra $\Lambda(s, s, 0)$ with $s \geq 1$ is called a *2-truncated cycle algebra*.

Now we observe the noncompact indecomposable objects in $\mathcal{D}^b(\Lambda(s, s, t))$. For $0 \leq p \leq s-1$, let X_p be the simple $\Lambda(s, s, t)$ -module S_p corresponding to the vertex p of the quiver $Q(s, t)$. For $-t \leq q \leq -1$, let Y_q be the indecomposable $\Lambda(s, s, t)$ -module P_q/S_1 if $s \geq 2$ and P_q/S_0 if $s = 1$, where P_q is the indecomposable projective $\Lambda(s, s, t)$ -module corresponding to the vertex q of the quiver $Q(s, t)$. Note that Y_q has a minimal projective resolution

$$\dots \xrightarrow{\alpha_0} P_0 \xrightarrow{\alpha_{s-1}} P_{s-1} \xrightarrow{\alpha_{s-2}} \dots \xrightarrow{\alpha_2} P_2 \xrightarrow{\alpha_1} P_1 \xrightarrow{\alpha_q \cdots \alpha_{-1} \alpha_0} P_q \longrightarrow 0$$

where all differentials are given by the left multiplications of the paths.

Lemma 1. *Let $A = \Lambda(s, s, t)$ with $s \geq 1$ and $t \geq 0$. Then, up to isomorphism and shift, $X_0, X_1, \dots, X_{s-1}$ and $Y_{-t}, Y_{-t+1}, \dots, Y_{-1}$ are all noncompact indecomposable objects in $\mathcal{D}^b(A)$. Moreover, for all noncompact indecomposable objects X and Y in $\mathcal{D}^b(A)$, the following statements hold:*

- (1) $\text{Hom}_{\mathcal{D}^b(A)}(X, X[h]) \cong k$ if $h = 0, s, 2s, \dots$, and $= 0$ otherwise.
- (2) $\text{RHom}_A(X, Y) \not\cong 0$.

Proof. For the gentle algebra $A = \Lambda(s, s, t)$, it is not difficult to see $GSt_c = \{\alpha_p, (\alpha_q \cdots \alpha_{-1} \alpha_0) \mid 0 \leq p \leq s-1, -t \leq q \leq -1\}$ and $GSt^c = \{\alpha_p^{-1}, (\alpha_0^{-1} \alpha_{-1}^{-1} \cdots \alpha_q^{-1}) \mid 0 \leq p \leq s-1, -t \leq q \leq -1\}$ (Ref. [5, 4.3]). Since generalized strings are defined up to inverse, i.e., the equivalence relation $\cong_s$ (Ref. [5, 4.1]), we have $GSt_c = GSt^c = \{\alpha_p, (\alpha_q \cdots \alpha_{-1} \alpha_0) \mid 0 \leq p \leq$

$s-1, -t \leq q \leq -1\}$. Due to [5, Definition 2, Lemma 6, and Theorem 3], up to isomorphism and shift, $X_0, X_1, \dots, X_{s-1}$ and $Y_{-t}, Y_{-t+1}, \dots, Y_{-1}$ are all noncompact indecomposable objects in $\mathcal{D}^b(A)$.

Let X and Y be two noncompact indecomposable objects in $\mathcal{D}^b(A)$.

(1) There are altogether two cases:

(1.1) $X = X_p, 0 \leq p \leq s-1$. Since X is the simple A -module S_p corresponding to the vertex p of the quiver $Q(s, t)$, it is easy to calculate $\text{Hom}_{\mathcal{D}A}(X, X[h]) \cong k$ if $h = 0, s, 2s, \dots$, and $= 0$ otherwise.

(1.2) $X = Y_q, -t \leq q \leq -1$. In the following, we only consider the case $s \geq 2$. For the case $s = 1$, it is enough to replace S_1 with S_0 in appropriate places. We have a short exact sequence $S_1 \rightarrowtail P_q \twoheadrightarrow Y_q$, which induces a triangle $P_q \rightarrow Y_q \rightarrow S_1[1] \rightarrow$ in $\mathcal{D}A$. Applying the derived functor $\text{RHom}_A(Y_q, -)$ to this triangle, we obtain a triangle $\text{RHom}_A(Y_q, P_q) \rightarrow \text{RHom}_A(Y_q, Y_q) \rightarrow \text{RHom}_A(Y_q, S_1[1]) \rightarrow$ in $\mathcal{D}k$. It is not difficult to prove $\text{RHom}_A(Y_q, P_q) \cong 0$. Thus we have $\text{RHom}_A(Y_q, Y_q) \cong \text{RHom}_A(Y_q, S_1[1])$. Now (1.2) follows from the obvious fact $H^h(\text{RHom}_A(Y_q, S_1[1])) \cong k$ if $h = 0, s, 2s, \dots$, and $= 0$ otherwise.

(2) There are altogether four cases:

(2.1) $(X = X_p) \wedge (Y = X_{p'}), 0 \leq p, p' \leq s-1$. Obviously, there exist infinitely many $h \in \mathbb{N}$ such that $\text{Ext}_A^h(S_p, S_{p'}) \neq 0$. Thus $\text{RHom}_A(X_p, X_{p'}) \not\cong 0$.

(2.2) $(X = X_p) \wedge (Y = Y_q), 0 \leq p \leq s-1, -t \leq q \leq -1$. Applying the derived functor $\text{RHom}_A(X_p, -)$ to the triangle $P_q \rightarrow Y_q \rightarrow S_1[1] \rightarrow$ in $\mathcal{D}A$, we obtain a triangle $\text{RHom}_A(X_p, P_q) \rightarrow \text{RHom}_A(X_p, Y_q) \rightarrow \text{RHom}_A(X_p, S_1[1]) \rightarrow$ in $\mathcal{D}k$ which gives a long exact sequence. We can show $\text{Hom}_{\mathcal{D}A}(X_p, P_q[h]) = 0$ for all $h \neq 0$. Thus $\text{Hom}_{\mathcal{D}A}(X_p, Y_q[h]) \cong \text{Hom}_{\mathcal{D}A}(X_p, S_1[h+1])$ for all $h \geq 1$. Since $\text{Ext}_A^h(S_p, S_1) \neq 0$ for infinitely many $h > 0$, there exist infinitely many $h \in \mathbb{N}$ such that $\text{Hom}_{\mathcal{D}A}(X_p, Y_q[h]) \neq 0$. Hence $\text{RHom}_A(X_p, Y_q) \not\cong 0$.

(2.3) $(X = Y_q) \wedge (Y = X_p), -t \leq q \leq -1, 0 \leq p \leq s-1$. Applying the derived functor $\text{RHom}_A(-, S_p)$ to the triangle $P_q \rightarrow Y_q \rightarrow S_1[1] \rightarrow$ in $\mathcal{D}A$, we obtain a triangle $\text{RHom}_A(S_1[1], S_p) \rightarrow \text{RHom}_A(Y_q, S_p) \rightarrow \text{RHom}_A(P_q, S_p) \rightarrow$ in $\mathcal{D}k$. Obviously, $\text{RHom}_A(P_q, S_p) \cong 0$. Therefore, we have $\text{RHom}_A(Y_q, S_p) \cong \text{RHom}_A(S_1[1], S_p) \not\cong 0$ by (2.1), i.e., $\text{RHom}_A(Y_q, X_p) \not\cong 0$.

(2.4) $(X = Y_q) \wedge (Y = Y_{q'}), -t \leq q, q' \leq -1$. Applying the derived functor $\text{RHom}_A(Y_q, -)$ to the triangle $P_{q'} \rightarrow Y_{q'} \rightarrow S_1[1] \rightarrow$ in $\mathcal{D}A$, we obtain a triangle $\text{RHom}_A(Y_q, P_{q'}) \rightarrow \text{RHom}_A(Y_q, Y_{q'}) \rightarrow \text{RHom}_A(Y_q, S_1[1]) \rightarrow$ in $\mathcal{D}k$. It is not difficult to show $\text{Hom}_{\mathcal{D}A}(Y_q, P_{q'}[h]) = 0$ for all $h \neq 1$. Thus $\text{Hom}_{\mathcal{D}A}(Y_q, Y_{q'}[h]) \cong \text{Hom}_{\mathcal{D}A}(Y_q, S_1[h+1])$ for all $h \geq 2$. Note that

$\text{Hom}_{\mathcal{D}A}(Y_q, S_1[h+1]) \neq 0$ for infinitely many $h \in \mathbb{N}$. Hence, there exist infinitely many $h \in \mathbb{N}$ such that $\text{Hom}_{\mathcal{D}A}(Y_q, Y_{q'}[h]) \neq 0$. Therefore, $\text{RHom}_A(Y_q, Y_{q'}) \not\cong 0$. $\square$

Before giving some results on the n -recollements of derived categories of derived discrete algebras, we recall the definition of n -recollement of triangulated categories.

Let $\mathcal{T}_1$, $\mathcal{T}$ and $\mathcal{T}_2$ be triangulated categories. A *recollement* of $\mathcal{T}$ relative to $\mathcal{T}_1$ and $\mathcal{T}_2$ is given by

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \mathcal{T}_1 & \xrightarrow{i_* = i_!} & \mathcal{T} & \xrightarrow{j^! = j^*} & \mathcal{T}_2 \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array}$$

such that

- (R1) (i^*, i_*) , $(i_!, i^!)$, $(j_!, j^!)$ and (j^*, j_*) are adjoint pairs of triangle functors;
- (R2) i_* , $j_!$ and j_* are full embeddings;
- (R3) $j^! i_* = 0$ (and thus also $i^! j_* = 0$ and $i^* j_! = 0$);
- (R4) for each $X \in \mathcal{T}$, there are triangles

$$\begin{aligned} j_! j^! X &\rightarrow X \rightarrow i_* i^* X \rightarrow \\ i_! i^! X &\rightarrow X \rightarrow j_* j^* X \rightarrow \end{aligned}$$

where the arrows to and from X are the counits and the units of the adjoint pairs respectively [6].

Let $\mathcal{T}_1$, $\mathcal{T}$ and $\mathcal{T}_2$ be triangulated categories, and n a positive integer. An n -*recollement* of $\mathcal{T}$ relative to $\mathcal{T}_1$ and $\mathcal{T}_2$ is given by $n+2$ layers of triangle functors

$$\begin{array}{ccccc} & \xleftarrow{\quad} & & \xleftarrow{\quad} & \\ \mathcal{T}_1 & \xrightarrow{\quad} & \mathcal{T} & \xrightarrow{\quad} & \mathcal{T}_2 \\ & \xleftarrow{\quad} & & \xleftarrow{\quad} & \\ & \xrightarrow{\quad} & & \xrightarrow{\quad} & \\ & \vdots & & \vdots & \end{array}$$

such that every consecutive three layers form a recollement [17].

From now on, we only focus on the n -recollements of derived categories of algebras, i.e., all three triangulated categories in an n -recollement are the derived categories of algebras.

Proposition 1. *Let A be a derived discrete algebra. Then every recollement of $\mathcal{D}A$ relative to $\mathcal{D}B$ and $\mathcal{D}C$ can be extended to an n -recollement for all $n \geq 1$.*

Proof. If $\text{gl.dim} A < \infty$ then this is clear by [3, Proposition 3.3]. If $\text{gl.dim} A = \infty$, all the connected directed summands of A that are of infinite global dimension are derived equivalent to $\Lambda(s, s, t)$ for some $s \geq 1$ and $t \geq 0$. Assume that there is a recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \mathcal{D}B & \xrightarrow{i_*} & \mathcal{D}A & \xrightarrow{j^!} & \mathcal{D}C \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array}.$$

Then $i_*B \in \mathcal{D}^b(A)$ by [17, Lemma 2]. If i_*B is not compact then there exists an indecomposable projective B -module P such that i_*P is not compact. Therefore, i_*P is a noncompact indecomposable object in $\mathcal{D}^b(\Lambda(s, s, t))$ for some $s \geq 1$ and $t \geq 0$. By Lemma 1, i_*P is isomorphic to a shift of X_p or Y_q , and $\text{Hom}_{\mathcal{D}A}(i_*P, i_*P[h]) \neq 0$ for infinitely many h . It is a contradiction, since i_*B , and thus i_*P , is exceptional. Hence, i_*B is compact and the recollement can be extended to a 2-recollement by [17, Lemma 3]. Furthermore, the recollement can be extended to an n -recollement for all $n \geq 1$ by induction. $\square$

The following result is a generalization of [33, Proposition], which states that derived discrete algebras are invariant under derived equivalence.

Proposition 2. *Let A , B and C be algebras, and $\mathcal{D}A$ admit a recollement relative to $\mathcal{D}B$ and $\mathcal{D}C$. If A is derived discrete then so are B and C .*

Proof. According to Proposition 1 and [17, Lemma 2], we have a recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \mathcal{D}B & \xrightarrow{i_*} & \mathcal{D}A & \xrightarrow{j^!} & \mathcal{D}C \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array},$$

where all six triangle functors can be restricted to $\mathcal{D}^b(\text{mod})$. Since i_* is fully faithful, it induces a map from the set of isomorphism classes of indecomposable objects in $\mathcal{D}^b(B)$ to the set of isomorphism classes of indecomposable objects in $\mathcal{D}^b(A)$. Due to $i^*i_* \cong 1$, this map is injective. Moreover, by the same estimate on the cohomological dimension vector as [33, Section 1.1], we can show that the cohomological dimension vectors of the images of indecomposable objects in $\mathcal{D}^b(B)$ of a fixed cohomological dimension vector under the triangle functor i_* are “bounded”. Thus B is derived discrete. Similarly, we can prove C is derived discrete by considering the triangle functor j_* instead of i_* . $\square$

Remark 1. The converse of Proposition 2 is not true in general. Kronecker algebra A , which is nothing but the path algebra kQ of the quiver Q given by two vertices 1 and 2 and two arrows from 1 to 2, provides a counterexample. Indeed, though $\mathcal{D}A$ admits a recollement relative to $\mathcal{D}k$ and $\mathcal{D}k$, A is not derived discrete since all derived discrete algebras are of finite representation type.

3 n -derived-simple derived discrete algebras

In this section, we will classify all n -derived-simple derived discrete algebras up to derived equivalence.

Let n be a positive integer. An algebra A is said to be n -derived-simple if its derived category $\mathcal{D}A$ admits no nontrivial n -recollements of derived categories of algebras [17]. Note that 1- (resp. 2-, 3-) derived-simple algebras here are just the $\mathcal{D}(\text{Mod})$ - (resp. $\mathcal{D}^-(\text{Mod})$ - or $\mathcal{D}^+(\text{Mod})$ -, $\mathcal{D}^b(\text{mod})$ - or $\mathcal{D}^b(\text{Mod})$ -) derived-simple algebras in [3].

Now we focus on the algebra $\Lambda(s, s, t)$, where $s \geq 1$ and $t \geq 0$. The following lemma shows that $\Lambda(s, s, t)$ is rigid when deconstructing its derived category by recollements.

Lemma 2. *Let $A = \Lambda(s, s, t)$ with $s \geq 1$ and $t \geq 0$, and $\mathcal{D}A$ admit a recollement relative to $\mathcal{D}B$ and $\mathcal{D}C$. Then either $\text{gl.dim}C < \infty$ and B is derived equivalent to $\Lambda(s, s, t') \oplus B'$ with $0 \leq t' \leq t$ and $\text{gl.dim}B' < \infty$, or $\text{gl.dim}B < \infty$ and C is derived equivalent to $\Lambda(s, s, t') \oplus C'$ with $0 \leq t' \leq t$ and $\text{gl.dim}C' < \infty$.*

Proof. If $\mathcal{D}A$ admits a recollement relative to $\mathcal{D}B$ and $\mathcal{D}C$, then both B and C are derived discrete by Proposition 2. Since $\text{gl.dim}A = \infty$, at least one of B and C is of infinite global dimension [3, Proposition 2.14]. Without loss of generality, we assume $\text{gl.dim}B = \infty$. Thus B is derived equivalent to $\tilde{B} = \Lambda(s', s', t') \oplus B'$ for some integers $s' \geq 1$ and $t' \geq 0$ and some algebra B' . Therefore, there is a recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \mathcal{D}\tilde{B} & \xrightarrow{i_*} & \mathcal{D}A & \xrightarrow{j^!} & \mathcal{D}C \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array}.$$

The proposition will be proved by the following three steps.

Step 1. We claim $s' = s$ and $t' \leq t$.

For this, let S'_0 be the simple $\Lambda(s', s', t')$ -module corresponding to the vertex 0 of the quiver $Q(s', t')$. Then S'_0 is naturally a $\tilde{B}$ -module as well and $\text{Hom}_{\mathcal{D}\tilde{B}}(S'_0, S'_0[h]) \cong k$ if $h = 0, s', 2s', \dots$, and $= 0$ otherwise. Since S'_0 is a noncompact indecomposable object in $\mathcal{D}^b(\tilde{B})$ and i^* preserves compactness, we have $i_*S'_0$ is a noncompact indecomposable object in $\mathcal{D}^b(A)$. By Lemma 1, we have $\text{Hom}_{\mathcal{D}A}(i_*S'_0, i_*S'_0[h]) \cong k$ if $h = 0, s, 2s, \dots$, and $= 0$ otherwise. Therefore, $s' = s$, since the triangle functor i_* is fully faithful. It follows from [3, Proposition 6.5] that the rank $r(\tilde{B})$ of the Grothendieck group $K_0(\tilde{B})$ of $\tilde{B}$ is not larger than the rank $r(A)$ of the Grothendieck group $K_0(A)$ of A . Hence, $t' \leq t$.

Step 2. We claim $\text{gl.dim} B' < \infty$.

Assume on the contrary $\text{gl.dim} B' = \infty$. Then B' is derived discrete by Proposition 2 and of infinite global dimension. Thus B' is derived equivalent to $\Lambda(s'', s'', t'') \oplus B''$ for some integers $s'' \geq 1$ and $t'' \geq 0$ and some algebra B'' . Therefore, i_* induces a triangle functor

$$i'_* : \mathcal{D}(\Lambda(s', s', t')) \oplus \mathcal{D}(\Lambda(s'', s'', t'')) \oplus \mathcal{D}B'' \simeq \mathcal{D}\tilde{B} \rightarrow \mathcal{D}A.$$

Let S''_0 be the simple $\Lambda(s'', s'', t'')$ -module corresponding to the vertex 0 of the quiver $Q(s'', t'')$. Then $\text{RHom}_A(i'_*S'_0, i'_*S''_0) \cong 0$. However, $i'_*S'_0$ and $i'_*S''_0$ are pairwise non-isomorphic noncompact indecomposable objects in $\mathcal{D}^b(A)$, and thus we have $\text{RHom}_A(i'_*S'_0, i'_*S''_0) \not\cong 0$ by Lemma 1. It is a contradiction.

Step 3. We claim $\text{gl.dim} C < \infty$.

Assume on the contrary $\text{gl.dim} C = \infty$. Then C is derived discrete by Proposition 2 and of infinite global dimension. Thus C is derived equivalent to $\tilde{C} = \Lambda(s''', s''', t''') \oplus C'$ for some integers $s''' \geq 1$ and $t''' \geq 0$ and some algebra C' . Therefore, there is a recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j!} & \\ \mathcal{D}\tilde{B} & \xrightarrow{i_*} & \mathcal{D}A & \xrightarrow{j^!} & \mathcal{D}\tilde{C} \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array}$$

where both i_* and j_* can be restricted to $\mathcal{D}^b(\text{mod})$ by Proposition 1 and [17, Lemma 2]. Let S'''_0 be the simple $\Lambda(s''', s''', t''')$ -module corresponding to the vertex 0 of the quiver $Q(s''', t''')$. Then $\text{RHom}_A(i_*S'_0, j_*S'''_0) \cong 0$. However, $i_*S'_0$ and $j_*S'''_0$ are noncompact indecomposable objects in $\mathcal{D}^b(A)$, and thus $\text{RHom}_A(i_*S'_0, j_*S'''_0) \not\cong 0$ by Lemma 1 (2). It is a contradiction. $\square$

As a consequence of Lemma 2, we have the following proposition which was obtained independently by Angeleri Hügel, König, Liu and Yang in different way in the earlier version of [3].

Proposition 3. *Every 2-truncated cycle algebra is n -derived-simple for all $n \geq 1$.*

Proof. Assume $A = \Lambda(s, s, 0)$ and $\mathcal{D}A$ admits an n -recollement relative to $\mathcal{D}B$ and $\mathcal{D}C$. According to Lemma 2, we may assume that $\text{gl.dim} C < \infty$ and B is derived equivalent to $\Lambda(s, s, 0) \oplus B'$ for some algebra B' . By [3, Proposition 6.5], we have $r(A) = r(B) + r(C) = r(A) + r(B') + r(C)$. Thus $r(B') = 0 = r(C)$. Hence $B' = 0 = C$, i.e., the n -recollement is trivial. $\square$

The next proposition implies that dropping an indecomposable projective module with projective radical in [24] induces a recollement of derived categories of algebras.

Proposition 4. *Let A be a basic algebra, e a primitive idempotent, $\bar{e} := 1 - e$, and $\text{rade}A$ a projective A -module. Then $\mathcal{D}A$ admits a recollement relative to $\mathcal{D}(\bar{e}A\bar{e})$ and $\mathcal{D}k$.*

Proof. Let X be the projective A -module $\bar{e}A$, and Y the simple A -module $eA/\text{rade}A$. Since $\text{rade}A$ is projective, we have $\text{pd}_A Y \leq 1$. It is easy to see that both X and Y are compact exceptional objects in $\mathcal{D}A$ with $\text{End}_A(X) \cong \bar{e}A\bar{e}$ and $\text{End}_A(Y) \cong k$, and $\text{RHom}_A(X, Y) \cong 0$. Moreover, for all $Z \in \mathcal{D}A$ satisfying $\text{RHom}_A(X, Z) \cong 0 \cong \text{RHom}_A(Y, Z)$, we have $\text{RHom}_A(X \oplus Y, Z) \cong 0$. Clearly, $\text{rade}A \in \text{thick}X$, the smallest thick subcategory of $\mathcal{D}A$ containing X , i.e., the smallest full triangulated subcategory of $\mathcal{D}A$ containing X and closed under direct summands. Thus eA , and further $A \cong eA \oplus X$, belongs to $\text{thick}(X \oplus Y)$. Hence $Z \cong \text{RHom}_A(A, Z) \cong 0$. Therefore, X and Y determine a recollement of $\mathcal{D}A$ relative to $\mathcal{D}k$ and $\mathcal{D}(\bar{e}A\bar{e})$ by [16, Proposition 1], and further a 2-recollement of $\mathcal{D}A$ relative to $\mathcal{D}k$ and $\mathcal{D}(\bar{e}A\bar{e})$ by [17, Proposition 1]. Hence, $\mathcal{D}A$ admits a recollement relative to $\mathcal{D}(\bar{e}A\bar{e})$ and $\mathcal{D}k$. $\square$

Let B be a bound quiver algebra, M a right B -module and N a left B -module. Then $\begin{bmatrix} B & 0 \\ M & k \end{bmatrix}$ is called a *one-point extension* of B , and $\begin{bmatrix} k & 0 \\ N & B \end{bmatrix}$ is called a *one-point coextension* of B (Ref. [30]). The following lemma implies that one-point (co)extension induces a recollement, which will be used frequently later.

Lemma 3. *Let A be a one-point (co)extension of a bound quiver algebra B . Then $\mathcal{D}A$ admits a recollement relative to $\mathcal{D}B$ and $\mathcal{D}k$.*

Proof. Both one-point extensions and one-point coextensions are special triangular matrix algebras. So this lemma follows from [17, Example 1 (2)]. In fact, the one-point coextension case also can be deduced from Proposition 4 by considering the idempotent element corresponding to the coextension vertex which is a sink in the quiver of A and corresponds to a simple projective A -module. $\square$

The following result is our main theorem in this section.

Theorem 1. *For any positive integer n , a derived discrete algebra is n -derived-simple if and only if it is derived equivalent to either k or a 2-truncated cycle algebra.*

Proof. Sufficiency. It follows from Proposition 3.

Necessity. Let A be a basic n -derived-simple derived discrete algebra. Then A must be connected. If A is a piecewise hereditary algebra of Dynkin type then it is derived equivalent to k by [2, Theorem 5.7]. Otherwise, A is derived equivalent to $\Lambda(r, s, t)$ for some $s \geq r \geq 1$ and $t \geq 0$. If $t \geq 1$ then A is a one-point extension of $\Lambda(r, s, t-1)$. It follows from Lemma 3 and Proposition 1 that $\mathcal{D}A$ admits a non-trivial n -recollement relative to $\mathcal{D}\Lambda(r, s, t-1)$ and $\mathcal{D}k$, which contradicts to the assumption. Hence, we have $t = 0$. If $s > r$ then applying Proposition 4 to the algebra $\Lambda(r, s, 0)$ and the idempotent e corresponding to the vertex 0 of the quiver $Q(s, 0)$ and also Proposition 1, we can obtain a non-trivial n -recollement of $\mathcal{D}\Lambda(r, s, 0)$, and thus $\mathcal{D}A$. It contradicts to the assumption, which implies $s = r$ and A must be derived equivalent to a 2-truncated cycle algebra $\Lambda(s, s, 0)$. $\square$

4 Jordan-Hölder theorems

In this section, we will show that the Jordan-Hölder theorems hold for all kinds of derived categories of derived discrete algebras.

Definition 1. Let n be a positive integer. An n -composition-series of the derived category $\mathcal{D}A$ of an algebra A is a chain of derived categories of algebras linked by fully faithful triangle functors

$$0 = \mathcal{D}B_l \xrightarrow{i_l} \mathcal{D}B_{l-1} \xrightarrow{i_{l-1}} \cdots \xrightarrow{i_2} \mathcal{D}B_1 \xrightarrow{i_1} \mathcal{D}B_0 = \mathcal{D}A$$

such that, for all $p = 1, 2, \dots, l$, the triangle functor $\mathcal{D}B_p \xrightarrow{i_p} \mathcal{D}B_{p-1}$ can be completed to an n -recollement

$$\begin{array}{ccccc} & \longleftarrow & & \longleftarrow & \\ \mathcal{D}B_p & \xrightarrow{i_p} & \mathcal{D}B_{p-1} & \longrightarrow & \mathcal{D}C_p \\ & \longleftarrow & & \longleftarrow & \\ & \vdots & & \vdots & \end{array}$$

for some n -derived-simple algebra C_p . In this case, $\mathcal{D}C_1, \mathcal{D}C_2, \dots, \mathcal{D}C_l$ are called the n -composition-factors of $\mathcal{D}A$. Moreover, $\mathcal{D}C_p$ is said to be of *multiplicity* m_p if it appears exactly m_p times in the sequence $\mathcal{D}C_1, \mathcal{D}C_2, \dots, \mathcal{D}C_l$ up to derived equivalence.

Remark 2. (1) By [3, Proposition 6.5], we know that the length l of an n -composition-series of $\mathcal{D}A$ is not larger than the rank $r(A)$ of the Grothendieck group $K_0(A)$ of A .

(2) If $n \neq n'$ then it is possible that the length of an n -composition-series of $\mathcal{D}A$ is not equal to the length of an n' -composition-series of $\mathcal{D}A$. For example, let A be the two-point algebra given in [3, Example 5.8] which is 2-derived simple but not 1-derived-simple. Then the length of any 1-composition-series of $\mathcal{D}A$ is 2. However, the length of any 2-composition-series of $\mathcal{D}A$ is 1.

(3) It is possible that the length of one n -composition-series of $\mathcal{D}A$ is equal to the length of the other n -composition-series of $\mathcal{D}A$ but two n -composition-series have different n -composition-factors, so the Jordan-Hölder theorems do not hold for derived categories of algebras in general. For example, let A be the two-point algebra given in [3, Example 7.6]. Then $\mathcal{D}A$ admits two 1-composition-series with completely different 1-composition-factors.

(4) According to [2, Theorem 5.7 and Corollary 5.9], for any piecewise hereditary algebra A and $n = 1, 2, 3$, $\mathcal{D}A$ admits an n -composition-series with only n -composition-factors $\mathcal{D}k$ of multiplicity $r(A)$ which are independent of the choice of n -composition-series up to derived equivalence. Thus the Jordan-Hölder theorem holds for the unbounded (resp. upper bounded, lower bounded, bounded) derived category of any piecewise hereditary algebra.

Lemma 4. Let $A_1, \dots, A_m$ be algebras, $A = \bigoplus_{p=1}^m A_p$ and n a positive integer. If $\mathcal{D}A_p$ admits an n -composition-series with n -composition-factors $\mathcal{D}C_{p,1}, \dots, \mathcal{D}C_{p,l_p}$ for all $1 \leq p \leq m$ then $\mathcal{D}A$ admits an n -composition-series with n -composition-factors $\mathcal{D}C_{1,1}, \dots, \mathcal{D}C_{1,l_1}, \dots, \mathcal{D}C_{m,1}, \dots, \mathcal{D}C_{m,l_m}$.

Proof. It is clear that any n -recollement

$$\begin{array}{ccccc} \mathcal{D}B & \xrightleftharpoons{\quad} & \mathcal{D}A & \xrightleftharpoons{\quad} & \mathcal{D}C \\ & \vdots & & \vdots & \end{array}$$

can always induce an n -recollement

$$\begin{array}{ccccc} \mathcal{D}(B \oplus E) & \xrightleftharpoons{\quad} & \mathcal{D}(A \oplus E) & \xrightleftharpoons{\quad} & \mathcal{D}C \\ & \vdots & & \vdots & \end{array}$$

for all algebras A, B, C and E . Let

$$0 = \mathcal{D}B_{p,l_p} \hookrightarrow \cdots \hookrightarrow \mathcal{D}B_{p,1} \hookrightarrow \mathcal{D}B_{p,0} = \mathcal{D}A_p$$

be an n -composition-series of $\mathcal{D}A_p$ with n -composition-factors $\mathcal{D}C_{p,1}, \dots, \mathcal{D}C_{p,l_p} = \mathcal{D}B_{p,l_p-1}$. Then they induce an n -composition-series $0 = \mathcal{D}B_{m,l_m} \hookrightarrow \cdots \hookrightarrow \mathcal{D}B_{m,1} \hookrightarrow \mathcal{D}B_{m,0} = \mathcal{D}A_m = \mathcal{D}A_m \oplus \mathcal{D}B_{m-1,l_{m-1}} \hookrightarrow \mathcal{D}A_m \oplus \mathcal{D}B_{m-1,l_{m-1}-1} \hookrightarrow \cdots \hookrightarrow \mathcal{D}A_m \oplus \mathcal{D}B_{m-1,1} \hookrightarrow \mathcal{D}A_m \oplus \mathcal{D}B_{m-1,0} = \mathcal{D}A_m \oplus \mathcal{D}A_{m-1} \hookrightarrow \cdots \hookrightarrow \mathcal{D}A$ of $\mathcal{D}A$ with n -composition-factors $\mathcal{D}C_{1,1}, \dots, \mathcal{D}C_{1,l_1}, \dots, \mathcal{D}C_{m,1}, \dots, \mathcal{D}C_{m,l_m}$. $\square$

The following result is our main theorem in this section.

Theorem 2. *Let n be a positive integer and A a derived discrete algebra, say derived equivalent to $\bigoplus_{p=1}^m A_p$ where $m \geq 1$ and $A_p = \Lambda(s_p, s_p, t_p)$ for $s_p \geq 1, t_p \geq 0, 1 \leq p \leq u$ and $0 \leq u \leq m$; $A_p = \Lambda(r_p, s_p, t_p)$ for $s_p > r_p \geq 1, t_p \geq 0, u+1 \leq p \leq v$ and $u \leq v \leq m$; A_p is a basic connected piecewise hereditary algebra of Dynkin type for $v+1 \leq p \leq m$. Then $\mathcal{D}A$ admits an n -composition-series with n -composition-factors $\mathcal{D}(\Lambda(s_1, s_1, 0)), \dots, \mathcal{D}(\Lambda(s_u, s_u, 0))$, and $\mathcal{D}k$ of multiplicity $r(A) - \sum_{p=1}^u s_p$. Moreover, any n -composition-series of $\mathcal{D}A$ has precisely these n -composition-factors up to reordering and derived equivalence.*

Proof. For $v+1 \leq p \leq m$, A_p is a basic connected piecewise hereditary algebra of Dynkin type. By [23, Theorem 1.1 (i)], A_p is *triangular*, i.e., its quiver has no oriented cycles. Thus A_p can be constructed from k by $r(A_p) - 1$ times of one-point extensions. It follows from Lemma 3, Proposition 2 and Proposition 1 that $\mathcal{D}A_p$ admits an n -composition-series with n -composition-factors $\mathcal{D}k$ of multiplicity $r(A_p)$.

For $u+1 \leq p \leq v$, $A_p = \Lambda(r_p, s_p, t_p)$ with $s_p > r_p \geq 1$ and $t_p \geq 0$. The algebra $\Lambda(r_p, s_p, t_p)$ can be constructed from $\Lambda(r_p, s_p, 0)$ by t_p times one-point extensions. For $\Lambda(r_p, s_p, 0)$, since $s_p > r_p$, applying Proposition 4 to the

idempotent element e corresponding to the vertex 0 of the quiver $Q(s_p, 0)$ and also Proposition 1, we get an n -recollement of $\mathcal{D}\Lambda(r_p, s_p, 0)$ relative to $\mathcal{D}E_p$ and $\mathcal{D}k$ where E_p is a Nakayama algebra whose quiver is a line quiver. The algebra E_p can be constructed from k by $r(E_p) - 1 = s_p - 2$ times of one-point extensions. Therefore, $\mathcal{D}\Lambda(r_p, s_p, t_p)$ admits an n -composition-series with n -composition-factors $\mathcal{D}k$ of multiplicity $r(\Lambda(r_p, s_p, t_p)) = t_p + s_p$.

For $1 \leq p \leq u$, $A_p = \Lambda(s_p, s_p, t_p)$ with $s_p \geq 1$ and $t_p \geq 0$. The algebra $\Lambda(s_p, s_p, t_p)$ can be constructed from $\Lambda(s_p, s_p, 0)$ by t_p times one-point extensions. Thus $\mathcal{D}\Lambda(s_p, s_p, t_p)$ admits an n -composition-series with n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ and $\mathcal{D}k$ of multiplicity t_p .

By the above analyzes and Lemma 4, we know $\mathcal{D}A$ admits an n -composition-series with n -composition-factors $\mathcal{D}(\Lambda(s_1, s_1, 0)), \dots, \mathcal{D}(\Lambda(s_u, s_u, 0))$, and $\mathcal{D}k$ of multiplicity $r(A) - \sum_{p=1}^u s_p$.

Next we prove these n -composition-factors of $\mathcal{D}A$ are independent of the choice of n -composition-series by induction on $r(A)$. If $r(A) = 1$ then A is local. Thus A is n -derived-simple. By Theorem 1, A is derived equivalent to either k or $\Lambda(1, 1, 0)$. In either case, we have nothing to say. Assume $r(A) \geq 2$ and the statement holds for all algebras B with $r(B) < r(A)$. For any n -composition-series

$$(*) \quad 0 = \mathcal{D}B_l \hookrightarrow \mathcal{D}B_{l-1} \hookrightarrow \dots \hookrightarrow \mathcal{D}B_1 \hookrightarrow \mathcal{D}B_0 = \mathcal{D}A$$

of $\mathcal{D}A$, we assume that the triangle functor $\mathcal{D}B_1 \hookrightarrow \mathcal{D}A$ is completed to an n -recollement

$$\begin{array}{ccccc} \mathcal{D}B_1 & \xrightleftharpoons{\quad} & \mathcal{D}A & \xrightleftharpoons{\quad} & \mathcal{D}C_1 \\ & \vdots & & \vdots & \end{array}$$

where C_1 is n -derived-simple. Then C_1 is derived discrete by Proposition 2. Thus up to derived equivalence $C_1 = k$ or $\Lambda(s, s, 0)$ for some integer $s \geq 1$ by Theorem 1.

According to the proof of [27, Corollary 3.4], we get $B_1 = \oplus_{p=1}^m B_{1,p}$, $C_1 = \oplus_{p=1}^m C_{1,p}$ and the above n -recollement can be decomposed as the direct sum of n -recollements

$$\begin{array}{ccccc} \mathcal{D}B_{1,p} & \xrightleftharpoons{\quad} & \mathcal{D}A_p & \xrightleftharpoons{\quad} & \mathcal{D}C_{1,p} \\ & \vdots & & \vdots & \end{array}$$

The n -derived-simplicity of C_1 implies that C_1 is connected. Therefore, there exists some $q \in \{1, 2, \dots, m\}$ such that $C_1 = C_{1,q}$ and $\mathcal{D}B_{1,p} \simeq \mathcal{D}A_p$ for all $p \neq q$. Up to derived equivalence, we may assume $B_1 = (\oplus_{p \neq q} A_p) \oplus B_{1,q}$.

If $1 \leq q \leq u$ then $A_q = \Lambda(s_q, s_q, t_q)$. By Lemma 2, we have either $C_1 = k$ and $B_{1,q} = \Lambda(s_q, s_q, t'_q) \oplus B'_{1,q}$ for some integer $0 \leq t'_q \leq t_q$ and some algebra $B'_{1,q}$ with $\text{gl.dim} B'_{1,q} < \infty$, or $C_1 = \Lambda(s_q, s_q, 0)$ and $\text{gl.dim} B_{1,q} < \infty$. If $u+1 \leq q \leq m$ then $\text{gl.dim} A_q < \infty$. Thus $C_1 = k$ and $\text{gl.dim} B_{1,q} < \infty$. Let's discuss these three cases in detail.

Case 1. $A_q = \Lambda(s_q, s_q, t_q)$, $C_1 = k$ and $B_{1,q} = \Lambda(s_q, s_q, t'_q) \oplus B'_{1,q}$ with $\text{gl.dim} B'_{1,q} < \infty$.

In this case $B_1 = (\oplus_{p \neq q} A_p) \oplus B_{1,q} = (\oplus_{p \neq q} A_p) \oplus \Lambda(s_q, s_q, t'_q) \oplus B'_{1,q}$, and $r(B_1) = r(A) - 1$. By induction assumption, the n -composition-series

$$0 = \mathcal{D}B_l \hookrightarrow \mathcal{D}B_{l-1} \hookrightarrow \cdots \hookrightarrow \mathcal{D}B_2 \hookrightarrow \mathcal{D}B_1$$

of $\mathcal{D}B_1$ has exactly n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ with $1 \leq p \leq u$ and $\mathcal{D}k$ of multiplicity $r(B_1) - \sum_{p=1}^u s_p$ up to reordering and derived equivalence. Thus the n -composition-series (*) of $\mathcal{D}A$ has exactly n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ with $1 \leq p \leq u$ and $\mathcal{D}k$ of multiplicity $r(B_1) - \sum_{p=1}^u s_p + 1 = r(A) - \sum_{p=1}^u s_p$ up to reordering and derived equivalence.

Case 2. $A_q = \Lambda(s_q, s_q, t_q)$, $C_1 = \Lambda(s_q, s_q, 0)$ and $\text{gl.dim} B_{1,q} < \infty$.

In this case $B_1 = (\oplus_{p \neq q} A_p) \oplus B_{1,q}$ with $\text{gl.dim} B_{1,q} < \infty$, and $r(B_1) = r(A) - s_q$. By induction assumption, the n -composition-series

$$0 = \mathcal{D}B_l \hookrightarrow \mathcal{D}B_{l-1} \hookrightarrow \cdots \hookrightarrow \mathcal{D}B_2 \hookrightarrow \mathcal{D}B_1$$

of $\mathcal{D}B_1$ has exactly n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ with $1 \leq p \leq u$ and $p \neq q$, and $\mathcal{D}k$ of multiplicity $r(B_1) - \sum_{p \neq q} s_p$, up to reordering and derived equivalence. Thus the n -composition-series (*) of $\mathcal{D}A$ has exactly n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ with $1 \leq p \leq u$ and $\mathcal{D}k$ of multiplicity $r(B_1) - \sum_{p \neq q} s_p = r(A) - \sum_{p=1}^u s_p$ up to reordering and derived equivalence.

Case 3. $\text{gl.dim} A_q < \infty$, $C_1 = k$ and $\text{gl.dim} B_{1,q} < \infty$.

In this case $B_1 = (\oplus_{p \neq q} A_p) \oplus B_{1,q}$ with $\text{gl.dim} B_{1,q} < \infty$ and $r(B_1) = r(A) - 1$. By induction assumption, the n -composition-series

$$0 = \mathcal{D}B_l \hookrightarrow \mathcal{D}B_{l-1} \hookrightarrow \cdots \hookrightarrow \mathcal{D}B_2 \hookrightarrow \mathcal{D}B_1$$

of $\mathcal{D}B_1$ has exactly n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ with $1 \leq p \leq u$ and $\mathcal{D}k$ of multiplicity $r(B_1) - \sum_{p=1}^u s_p$ up to reordering and derived equivalence. Thus the n -composition-series (*) of $\mathcal{D}A$ has exactly n -composition-factors $\mathcal{D}(\Lambda(s_p, s_p, 0))$ with $1 \leq p \leq u$ and $\mathcal{D}k$ of multiplicity $r(B_1) - \sum_{p=1}^u s_p + 1 = r(A) - \sum_{p=1}^u s_p$ up to reordering and derived equivalence.

Now we finish the proof of the theorem. $\square$

Remark 3. For $n = 1$ (resp. 2, 3), Theorem 2 implies that the Jordan-Hölder theorem holds for the unbounded (resp. upper bounded or lower bounded, bounded) derived category of any derived discrete algebra.

ACKNOWLEDGMENT. The author is grateful to her supervisor Professor Yang Han for his guidance. This work is supported by Project 11171325 NSFC.

References

- [1] L. Angeleri Hügel, S. König and Q. Liu, On the uniqueness of stratifications of derived module categories, *J. Algebra* 359 (2012), 120–137.
- [2] L. Angeleri Hügel, S. König and Q. Liu, Jordan-Hölder theorems for derived module categories of piecewise hereditary algebras, *J. Algebra* 352 (2012), 361–381.
- [3] L. Angeleri Hügel, S. König, Q. Liu and D. Yang, Derived simple algebras and restrictions of recollements of derived module categories, arXiv:1310.3479v1 [math.RT].
- [4] I. Assem, D. Simson and A. Skowroński, Elements of the representation theory of associative algebras, 1 : Techniques of representation theory, London Mathematical Society Student Texts, vol. 65, Cambridge University Press, Cambridge, 2006.
- [5] V. Bekkert and H. Merklen, Indecomposables in derived categories of gentle algebras, *Alg. Rep. Theory* 6 (2003), 285–302.
- [6] A.A. Beilinson, J. Bernstein and P. Deligne, Faisceaux pervers, *Astérisque*, vol. 100, Soc. Math. France, 1982.
- [7] G. Bobiński, The graded centers of derived discrete algebras, *J. Algebra* 333 (2011), 55–66.
- [8] G. Bobiński, C. Geiß and A. Skowroński, Classification of discrete derived categories, *Central Eur. J. Math.* 2 (2004), 19–49.
- [9] G. Bobiński and H. Krause, The Krull-Gabriel dimension of discrete derived categories, *Bull. Sci. Math.* To appear.
- [10] M. Bökstedt and A. Neeman, Homotopy limits in triangulated categories, *Compositio Math.* 86 (1993), 209–234.
- [11] N. Broomhead, D. Pauksztello and D. Ploog, Discrete derived categories I: homomorphisms, autoequivalences and t -structures, arXiv:1312.5203v1 [math.RT].
- [12] N. Broomhead, D. Pauksztello and D. Ploog, Discrete derived categories II: The silting pairs CW complex and the stability manifold, arXiv:1407.5944v1 [math.RT].
- [13] H.X. Chen and C.C. Xi, Stratifications of derived categories from tilting modules over tame hereditary algebras, arXiv:1107.0444 [math.RT].
- [14] H.X. Chen and C.C. Xi, Good tilting modules and recollements of derived module categories, *Proc. Lond. Math. Soc.* 104 (2012), no. 5, 959–996.

- [15] H.X. Chen and C.C. Xi, Recollements of derived categories III: Finitistic dimensions, arXiv:1405.5090v1 [math.RA].
- [16] Y. Han, Recollements and Hochschild theory, *J. Algebra* 397 (2014), 535–547.
- [17] Y. Han and Y.Y. Qin, Reducing homological conjectures by n -recollements, arXiv:1410.3223v2 [math.RT].
- [18] Y. Han and C. Zhang, Brauer-Thrall type theorems for derived category, arXiv:1310.2777v1 [math.RT].
- [19] Z. Han, Homotopy categories of injective modules over derived discrete algebras, arXiv:1308.2334v1 [math.RT].
- [20] D. Happel, Triangulated categories in the representation theory of finite dimensional algebras, *London Math. Soc. Lecture Notes Ser.* 119, 1988.
- [21] D. Happel, A family of algebras with two simple modules and Fibonacci numbers, *Arch. Math. (Basel)* 57 (1991), no. 2, 133–139.
- [22] D. Happel, Reduction techniques for homological conjectures, *Tsukuba J. Math.* 17 (1993), 115–130.
- [23] D. Happel, I. Reiten, and S.O. Smalø, Piecewise hereditary algebras, *Arch. Math. (Basel)* 66 (1996), 182–186.
- [24] K. Igusa and D. Zacharia, On the cyclic homology of monomial relation algebras, *J. Algebra* 151 (1992), 502–521.
- [25] B. Keller, Invariance and localization for cyclic homology of DG algebras, *J. Pure Appl. Algebra* 123 (1998), no. 1–3, 223–273.
- [26] S. König, Tilting complexes, perpendicular categories and recollements of derived module categories of rings, *J. Pure Appl. Algebra* 73 (1991) 211–232.
- [27] Q. Liu and D. Yang, Blocks of group algebras are derived simple, *Math. Z.* 272 (2012), 913–920.
- [28] Q. Liu and D. Yang, Stratifications of algebras with two simple modules, *Forum Math.* To appear.
- [29] A. Neeman and A. Ranicki, Noncommutative localisation in algebraic K-theory I, *Geometry and topology* 8 (2004), 1385–1425.
- [30] C.M. Ringel, Tame algebras and integral quadratic forms, *Lecture Notes in Math.*, vol. 1099, Springer-Verlag, Berlin, 1984.
- [31] M. Schlichting, Negative K-theory of derived categories, *Math. Z.* 253 (2006), no. 1, 97–134.
- [32] R.W. Thomason and T. Trobaugh, Higher algebraic K-theory of schemes and of derived categories, in: *The Grothendieck Festschrift, III*, Birkhäuser, Progress Math. 87 (1990), 247–436.
- [33] D. Vossieck, The algebras with discrete derived category, *J. Algebra* 243 (2001), 168–176.

- [34] A. Wiedemann, On stratifications of derived module categories, *Canad. Math. Bull.* 34 (1991), 275–280.
- [35] D. Yang, Algebraic stratifications and derived simplicity, Plenary talk on ICRA XVI, 2014.
- [36] D. Yao, Higher algebraic K-theory of admissible abelian categories and localization theorems, *J. Pure Appl. Algebra* 77 (1992), 263–339.